

Fibonacci Identities via the Determinant Sum Property

Michael Z. Spivey



Mike Spivey (mspivey@ups.edu) received his B.S. from Samford University and his Ph.D. from Princeton, and he currently teaches at the University of Puget Sound (Tacoma, WA 98416). His research background is in optimization, but he also enjoys dabbling in combinatorics and elementary number theory. Lately, he hasn't been able to stop himself from playing around with matrices made up of various mathematical entities.

There is a long tradition of using matrices and determinants to study Fibonacci numbers. For example, Bicknell-Johnson and Spears [3] use elementary matrix operations and determinants to generate classes of identities for generalized Fibonacci numbers, and Cahill and Narayan [4] show how Fibonacci and Lucas numbers arise as determinants of some tridiagonal matrices. Another recent paper, by Benjamin, Cameron, and Quinn [2], provides combinatorial interpretations for Fibonacci identities using determinants. Koshy's extensive survey text [7] of the Fibonacci numbers includes two chapters on the use of matrices and determinants as well.

However, none of this work utilizes the sum property for determinants [6]:

If A , B , and C are matrices with identical entries except that one row (column) of C , say the k th, is the sum of the k th rows (columns) of A and B , then $|A| + |B| = |C|$.

In this note we use this property to give a new proof of the following identity for the Fibonacci numbers [1]:

$$F_m F_n - F_{m-r} F_{n+r} = (-1)^{m-r} F_r F_{n+r-m}. \tag{1}$$

The following identities [8] are special cases of (1):

Cassini's identity: $F_{n+1} F_{n-1} - F_n^2 = (-1)^n \tag{2}$

d'Ocagne's identity: $F_{m+1} F_n - F_m F_{n+1} = (-1)^m F_{n-m} \tag{3}$

Catalan's identity: $F_n^2 - F_{n+r} F_{n-r} = (-1)^{n-r} F_r^2 \tag{4}$

Our proof of (1) has three stages; we reach Cassini's identity at the end of the first stage and d'Ocagne's at the end of the second.

Generalized Fibonacci sequences

The Fibonacci sequence is defined by the initial values $F_0 = 0$ and $F_1 = 1$ and the recurrence relation $F_n = F_{n-1} + F_{n-2}$, where $n \geq 2$. A generalized Fibonacci sequence is a sequence of numbers $\{G_n\}$ that obeys the same Fibonacci recurrence relation but

with arbitrary starting values. Thus $G_n = G_{n-1} + G_{n-2}$, while G_0 and G_1 can be any numbers. It can readily be shown ([5], p. 493), that

$$G_n = G_0 F_{n-1} + G_1 F_n. \quad (5)$$

First stage of the proof

We begin with the 2×2 identity matrix:

$$A_0 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}.$$

For each $n \geq 1$, we now recursively construct A_n from A_{n-1} by adding the second row of A_{n-1} to the first and then interchanging the two rows.

Then, for example,

$$A_1 = \begin{bmatrix} 0 & 1 \\ 1 & 1 \end{bmatrix}, \quad A_2 = \begin{bmatrix} 1 & 1 \\ 1 & 2 \end{bmatrix}, \quad A_3 = \begin{bmatrix} 1 & 2 \\ 2 & 3 \end{bmatrix}, \quad \text{and} \quad A_4 = \begin{bmatrix} 2 & 3 \\ 3 & 5 \end{bmatrix}.$$

The entries in each A_n appear to be Fibonacci numbers, and, in fact, it is easy to show by induction that

$$A_n = \begin{bmatrix} F_{n-1} & F_n \\ F_n & F_{n+1} \end{bmatrix}.$$

The two steps involved in generating A_n from A_{n-1} are both elementary matrix row operations: In the first, one row is added to another, while in the second, two rows are swapped. The first of these does not affect the determinant, and the second changes only the sign. Since $|A_0| = 1$, it follows that $|A_n| = (-1)^n$. Since

$$A_n = \begin{bmatrix} F_{n-1} & F_n \\ F_n & F_{n+1} \end{bmatrix},$$

the definition of the determinant implies that $F_{n+1}F_{n-1} - F_n^2 = (-1)^n$, Cassini's identity (2). (We note that there is another proof of Cassini's identity that uses determinants; see, for example, Knuth [5], p. 81.)

Second stage

The second stage brings us to d'Ocagne's identity. Let B_0 be the matrix

$$\begin{bmatrix} F_n & F_n \\ F_{n+1} & F_{n+1} \end{bmatrix}.$$

Construct B_1 by adding column 1 of A_n to that of B_0 . Thus

$$B_1 = \begin{bmatrix} F_{n+1} & F_n \\ F_{n+2} & F_{n+1} \end{bmatrix},$$

and by the determinant sum property, $|B_1| = (-1)^n$. For $r \geq 2$, we now construct B_r by adding the first column of B_{r-2} to that of B_{r-1} . Thus,

$$B_2 = \begin{bmatrix} F_{n+2} & F_n \\ F_{n+3} & F_{n+1} \end{bmatrix}, B_3 = \begin{bmatrix} F_{n+3} & F_n \\ F_{n+4} & F_{n+1} \end{bmatrix}, B_4 = \begin{bmatrix} F_{n+4} & F_n \\ F_{n+5} & F_{n+1} \end{bmatrix},$$

and so forth.

An easy induction proof shows that

$$B_r = \begin{bmatrix} F_{n+r} & F_n \\ F_{n+r+1} & F_{n+1} \end{bmatrix}. \tag{6}$$

From the determinant sum property, we have $|B_r| = |B_{r-1}| + |B_{r-2}|$. This recurrence relation implies that the sequence $\{|B_r|\}$ is actually a generalized Fibonacci sequence. Since $|B_0| = 0$ and $|B_1| = (-1)^n$, it follows from (5) that for $r \geq 0$,

$$|B_r| = (-1)^n F_r. \tag{7}$$

Since $|B_0| = 0$, not only is the sequence $\{|B_r|\}$ a generalized Fibonacci sequence, it is actually the Fibonacci sequence or its negative.

By (6) and the definition of the determinant, $|B_r| = F_{n+r}F_{n+1} - F_nF_{n+r+1}$. Let $m = n + r$, and it follows from this result and (7) that $F_{n+1}F_m - F_nF_{m+1} = (-1)^n F_{m-n}$, d’Ocagne’s identity (3). Consequently, we can see that the Fibonacci numbers on the right-hand side of d’Ocagne’s identity arise naturally as a result of adding determinants.

Third stage

In the final stage of the proof we again generate a sequence of matrices, but instead of adding the first columns of the two previous matrices together to obtain the next matrix in the sequence, we add the bottom rows. Let

$$C_0 = \begin{bmatrix} F_n & F_{n-r} \\ F_n & F_{n-r} \end{bmatrix} \quad \text{and let} \quad C_1 = \begin{bmatrix} F_n & F_{n-r} \\ F_{n+1} & F_{n-r+1} \end{bmatrix},$$

the matrix obtained by replacing n with $n - r$ in B_r . Now, for $s \geq 2$, construct C_s by adding the second row of C_{s-2} to that of C_{s-1} .

Then

$$C_2 = \begin{bmatrix} F_n & F_{n-r} \\ F_{n+2} & F_{n-r+2} \end{bmatrix}, C_3 = \begin{bmatrix} F_n & F_{n-r} \\ F_{n+3} & F_{n-r+3} \end{bmatrix}, C_4 = \begin{bmatrix} F_n & F_{n-r} \\ F_{n+4} & F_{n-r+4} \end{bmatrix},$$

and so forth. Again, we can show by induction that

$$C_s = \begin{bmatrix} F_n & F_{n-r} \\ F_{n+s} & F_{n-r+s} \end{bmatrix}. \tag{8}$$

As in the second stage of the proof, it follows from the determinant sum property that $|C_s| = |C_{s-1}| + |C_{s-2}|$. This means that the sequence $\{|C_s|\}$ is also a generalized Fibonacci sequence. By (7), $|C_1| = (-1)^{n-r} F_r$. Since $|C_0| = 0$, it then follows from (5) that for $s \geq 0$,

$$|C_s| = [(-1)^{n-r} F_r] F_s. \tag{9}$$

Like the determinants $|B_r|$, the determinants $|C_s|$ form a multiple of the Fibonacci sequence.

By (8) and the definition of the determinant, $|C_s| = F_n F_{n-r+s} - F_{n+s} F_{n-r}$. Letting $s = m - n + r$, we see from this result and (9) that $F_n F_m - F_{m+r} F_{n-r} = (-1)^{n-r} F_r F_{m-n+r}$, which is (1). Catalan's identity (4) is the special case $m = n$. As in Stage 2 and the proof of d'Ocagne's identity, the Fibonacci numbers on the right-hand side of (1) arise naturally as a result of the generalized Fibonacci sequence of determinants obtained by the repeated use of the determinant sum property.

Acknowledgement. The author would like to thank an anonymous referee for many suggestions that improved the exposition in this paper.

References

1. S. Basin and V. Hoggatt, Jr., A primer on the Fibonacci sequence, Part II, *Fibonacci Quart.* **1** (1963) 61–68.
2. A. Benjamin, N. Cameron, and J. Quinn, Fibonacci determinants—a combinatorial approach, *Fibonacci Quart.*, to appear.
3. M. Bicknell-Johnson and C. Spears, Classes of identities for the generalized Fibonacci numbers $G_n = G_{n-1} + G_{n-c}$ from matrices with constant valued determinants, *Fibonacci Quart.* **34** (1996) 121–128.
4. N. Cahill and D. Narayan, Fibonacci and Lucas numbers as tridiagonal matrix determinants, *Fibonacci Quart.* **42** (2004) 216–221.
5. D. Knuth, *The Art of Computer Programming, Vol. I: Fundamental Algorithms*, 3rd ed., Addison-Wesley, 1997.
6. B. Kolman and D. Hill, *Elementary Linear Algebra*, 8th ed., Pearson, 2004.
7. T. Koshy, *Fibonacci and Lucas Numbers with Applications*, Wiley, 2001.
8. E. Weisstein et al., Fibonacci number, from *MathWorld*—A Wolfram Web Resource, <http://mathworld.wolfram.com/FibonacciNumber.html>.